

Postoperative Care

Introduction

The goal of postoperative care is to minimize complications of anesthesia and surgery by early detection and prompt treatment. After receiving anesthesia patients may experience pain, inadequate oxygenation, or adverse physiologic effects of sudden movement.

Recovery from general anesthesia takes longer than induction because the anesthetic is retained in fat and muscle. Fat has a meager blood supply; thus, it releases the anesthetic slowly, providing enough anesthesia to maintain adequate blood and brain levels during surgery. The patient's recovery time varies with his amount of body fat, his overall condition including renal function, his pre-medication regimen, and the type, dosage, and duration of anesthesia.

Physician's Orders

Check the physician's orders for:

- ❖ Pain management
- ❖ Other medication orders
- ❖ IV fluids/diet
- ❖ Wound/drain care
- ❖ Position/activity/C & DB-patients should ambulate as soon as possible
- ❖ Lab work or other tests
- ❖ Other orders

The PACU Record

Obtain and review the patient's PACU record, paying particular attention to:

- ❖ Summary of operative procedures and pertinent findings
- ❖ Type of anesthesia and any other medications given-preoperative, intraoperative, and postoperative
- ❖ Vital signs
- ❖ Medical history, including medications
- ❖ Fluid therapy, including estimated blood loss
- ❖ Type and number of drains, catheters and note amount and characteristics of any drainage
- ❖ The location(s) and condition of the surgical wound

The Postoperative Assessment

A systematic approach helps to ensure a thorough assessment:

- ❖ Overall appearance
- ❖ Level of consciousness, skin color, mucous membranes
- ❖ Respiratory status-breathing pattern, rate, depth, auscultate breath sounds; monitor O₂ saturation and obtain an order to administer oxygen if needed
- ❖ Pulse-rate and quality-should be within 20% of the pre-op heart rate
- ❖ Blood Pressure-should be within 20% of pre-op BP
- ❖ Temperature-anesthesia lowers body temperature; use blankets as needed
- ❖ Pain-intensity, characteristics, location, frequency, duration; provide medication as needed/ordered
- ❖ Other symptoms or sources of discomfort-nausea, vomiting, positioning, anxiety; provide medication as needed/ordered
- ❖ Infusion sites-assess for redness, swelling, infiltrates, pain, drainage
- ❖ Surgical wound dressings-assess drainage, amount, characteristics, odors
- ❖ Presence and condition of drains and tubes-color, type, odor, amount of drainage and urinary output. **Make sure all tubes and drains are properly connected and free of kinks and obstructions.**
- ❖ Post vascular or orthopedic surgery-assess extremity(s) for color, temperature, sensation, movement, presence and quality of pulses
- ❖ Antiembolic stockings/PAS stockings-apply to **non-surgical** leg(s)
- ❖ Monitor intake and output closely
- ❖ Assess for the presence of bowel sounds and passage of flatus before offering food to patients following abdominal surgery-check physician orders
- ❖ Encourage coughing and deep breathing exercises
- ❖ Patient controlled anesthesia-instruct patient on how to use (if being used)
- ❖ Instruct the patient on the use of the call bell
- ❖ Assess lab work-Lab values often change within the first 2 days following surgery as a result of blood and fluid loss and the body's reaction to surgery

Special considerations

- ❖ Fear, pain, anxiety, hypothermia, confusion, and immobility can upset the patient and jeopardize his safety and postoperative status. Offer emotional support to the patient and his family. Keep in mind that the patient who has lost a body part or who has been diagnosed with an incurable disease will need ongoing emotional support. Refer him and his family for counseling as needed.
- ❖ If the patient has **epidural analgesia infusion** for postoperative pain control, monitor his respiratory status closely. *Respiratory arrest may result from the respiratory depressant effects of the opioid.* Also monitor the patient's blood

pressure, heart rate, and arterial oxygen saturation. He may also suffer nausea, vomiting, or itching. Epidural analgesia may also include administering a local anesthetic with the opioid. Assess the patient's lower-extremity motor strength every 2 to 4 hours. If sensorimotor loss occurs (numbness or weakness of the legs), notify the anesthesiologist because the dosage may need to be decreased.

- ❖ If the patient is older, be aware of age-related changes that will alter your assessment. Cardiovascular status should be monitored carefully *because blood loss, pain, bed rest, and fluid and electrolyte imbalances can alter it*. Respiratory status also should be monitored carefully *because ventilation and oxygenation can be altered by age-related changes or years of smoking or chronic diseases*. Monitor level of consciousness and pain carefully *because mental status changes can affect these and make pain control more difficult*. Drug metabolism slows *with age* so monitor the older adult's risk for drug reactions, toxicity, and interactions. Monitor intake and output carefully and watch for urinary tract infections because of decreased renal functioning and decreased bladder capacity. Be careful when positioning the older patient and prevent postoperative falls *because the patient may have osteoporosis and may be prone to fractures*. Also monitor for signs and symptoms of infection *because risk for infection increases with age*.
- ❖ **All post-operative patients are considered a fall risk for the first 24 hours** following surgery. Assist patients with ADLs and mobility during this time. Assessment of fall risk is ongoing!

Complications

Postoperative complications may include arrhythmias, bleeding, hypotension, hypovolemia, septicemia, septic shock, atelectasis, pneumonia, thrombophlebitis, pulmonary embolism, urine retention, wound infection, wound dehiscence, evisceration, abdominal distention, paralytic ileus, constipation, altered body image, and postoperative psychosis.

Documentation

Document assessments and vital signs on the appropriate flowsheet. Record the condition of dressings, and drains, and characteristics of drainage. Document all interventions taken to alleviate pain and anxiety and the patient's responses to them. Document any complications and interventions taken, as well as patient response.

The Whipple procedure

Better outcomes for pancreatic cancer surgery.

Pancreatic cancer has been in the public eye lately because it has afflicted several prominent people, including Supreme Court Justice Ruth Bader Ginsburg, Apple CEO Steve Jobs, actor Patrick Swayze, and Randy Pausch, a computer science professor at Carnegie Mellon University whose inspirational last lecture became a YouTube sensation and, as a book, a national best seller.

It's odd that the disease would strike so many famous people at about the same time, because pancreatic cancer is fairly uncommon. Nearly 38,000 Americans were diagnosed with the disease last year, a fraction of the 215,000 who will be diagnosed with lung cancer. The media attention isn't hard to fathom, though. It's morbid interest: no other common cancer has such a poor prognosis. Only about 5% of those diagnosed with pancreatic cancer are alive five years later, in contrast to about 66% of colon cancer patients and 90% of female breast cancer patients. For a variety of reasons—screening, earlier diagnosis, better treatment—cancer isn't the proverbial death sentence it once was. For many, it is a survivable (if harrowing) condition that can be lived with for many years in relative good health. Pancreatic cancer stands out as a throwback.

But one bright spot in pancreatic cancer treatment is improved results for the Whipple procedure, the operation most often used to treat the disease. In the 1970s, over 15% of the patients who had the procedure died during the operation or shortly afterward. Improvements in surgical technique, anesthesia, and postoperative care have driven that rate into the low single digits at some hospitals, and the five-year survival rate after the operation may be as high as 20%. Studies have consistently shown that results are better at hospitals where many Whipples are done, and the operation is held up as an example of why steering patients

to high-volume centers for complex surgeries and treatments might be one way to improve the quality of health care and treatment outcomes.

One head, two hats

A healthy pancreas is spongy, yellowish-tan, and about seven inches long. It has a creaturely shape, with a large head that nestles into a loop of the first part of the small intestine (the duodenum), a longish body that squeezes behind the stomach, and a narrower tail that reaches the spleen (see the illustration on the next page). Physiologically, the pancreas wears two hats. It contains exocrine tissue that produces digestive enzymes, which are transported via ducts to the small intestine, and endocrine tissue that produces hormones, including insulin and glucagon. Over 90% of pancreatic cancers come from the exocrine part, and most of them are ductal adenocarcinomas—cancers that form in the lining of the organ's elaborate ductwork.

Some pancreatic cancers are caught early, discovered incidentally on computed tomography (CT) scans and other imaging studies ordered for unrelated reasons. That's apparently what happened in Ginsburg's case.

But for the most part, pancreatic cancer is diagnosed after someone has symptoms, which typically include abdominal pain, weight loss (common with cancer but especially so with pancreatic cancer), and jaundice, a yellowing of the skin and the whites of the eyes from a buildup of bilirubin in the blood, which can occur when a pancreatic tumor impinges on the common bile duct. A major reason pancreatic cancer is so lethal is that the cancer grows and spreads long before it causes any symptoms.

Surgical candidates

At the time of diagnosis, about 40% of pancreatic cancer patients have cancer that has already spread (metastasized)

extensively outside the organ. Surgery isn't an option once that has happened. Another 40% of patients have "locally advanced disease": the cancer hasn't metastasized, but it may have adhered to or invaded adjacent structures. The pancreas wraps around two large blood vessels, the superior mesenteric vein and artery. If the cancer gets intertwined with those blood vessels, that may preclude surgery.

Locally advanced pancreatic cancer can be treated with radiation and chemotherapy, but the median survival time is eight to 12 months. It's even shorter for people whose cancers have metastasized.

That leaves about 20% of pancreatic cancer patients with tumors that are, in the words of cancer specialists, "resectable"—that is, they can be treated surgically. Most of these tumors are confined to the head of the pancreas or its extension, the uncinate process, and the Whipple procedure is the preferred operation. By the time the cancer is in the body or tail of the pancreas, it's usually too late to operate, although that's not always the case.

The Whipple procedure

Pancreaticoduodenectomy, the formal name for the Whipple procedure, is a mouthful, so even doctors prefer the eponym. The procedure is named for Dr. Allen O. Whipple, the first American surgeon to perform the operation in 1935.

Because resectable pancreatic cancer is limited to the head of the pancreas in most cases, you might think that the operation to remove it would involve taking out just that part of the pancreas. But the head of the pancreas is structurally tied in to other organs and ducts, and it shares a common blood supply with them, so to remove it requires a much more extensive operation, namely the Whipple. A Whipple involves removing the head of the pancreas, the duodenum, the common bile duct, the

gallbladder, and often part of the stomach (see illustration). Surgeons then seal off the end of the small intestine and reattach what's left of the bile duct, pancreas (body and tail), and stomach.

Patients typically spend a week in the hospital. The recovery at home can be slow and fairly painful, so painkillers of some kind are usually needed. Initially, Whipple patients can eat only very small amounts of food that are very easy to digest, and they may need to take pancreatic enzymes to help with digestion, particularly of fatty foods. Diarrhea can be a problem that makes getting out of the house difficult. But, remarkably, the rearranged and plumbed digestive tract manages to recover in two to three months. "Last night I had steak and steak fries, and I'm doing pretty well," a 37-year-old patient told us in early February. He had his Whipple operation in October 2008.

One variation on the Whipple keeps the stomach intact. Pylorus-preserving Whipples (the pylorus is the muscular opening of the stomach that attaches to the duodenum) reduce surgical time and, theoretically, improve digestion and nutrition. Surgeons at Johns Hopkins Hospital favor the pylorus-preserving procedure, but Dr. Carlos Fernández-del Castillo, a surgeon at Harvard-affiliated Massachusetts General Hospital, said surgeons there don't do them for two reasons. First, studies haven't shown any long-term benefit. Second, patients are more likely to need intravenous feeding and stay in the hospital longer.

The most common complication immediately after surgery is leakage of pancreatic juices from the remnant of the pancreas. This may be more of a problem for patients with healthier,

more productive pancreases. Leaks can be treated with slender drains that channel the juices out of the body so they don't collect inside the abdomen. Dr. Fernández-del Castillo says a CT scan can help doctors decide if a leak is serious and needs treatment.

cited study published in *The New England Journal of Medicine* several years ago, the death rate at low-volume hospitals (those where less than one Whipple a year was performed, on average) was four times higher (16.3% vs. 3.8%) than at high-volume hospitals (more than 16 procedures a year).

The same high-volume-better-outcome math seems to apply to surgeons. According to one study, operative mortality rates varied by a factor of nearly four depending on the number of Whipples performed by the surgeon, even at a high-volume hospital.


Still an uphill battle

The post-Whipple prognosis is brightest for patients whose cancers have not spread to nearby lymph nodes. For these "node negative" patients, the five-year survival rate is 25% to 30%. For node-positive patients, it's only about 10%. Regardless of their lymph node status, most Whipple patients will get radiation therapy, chemotherapy, or both, to improve these odds, but cancer specialists haven't settled on the right combination nor precisely which drugs should be used (there have been encouraging results for a drug called gemcitabine).

But another sobering way to view those survival percentages is to consider the larger percentage of Whipple patients who don't make it to the five-year mark. Randy Pausch is an example. He was

diagnosed in the summer of 2006, had a Whipple and follow-up radiation and chemotherapy. Pausch was 47 when he died in July 2008. The improvements in the Whipple operation are a welcome development, but it's a relatively small step in the long, uphill battle against pancreatic cancer. ♥

The Whipple procedure for pancreatic cancer



More is better

Many studies have shown that the outcomes for an operation tend to be better at hospitals where those operations are performed often. This commonsensical relationship seems to be especially true of the Whipple procedure, probably because of its complexity. In one frequently

Source: from Harvard Health Letter, Harvard Health Publications, Copyright 2009 by President and Fellows of Harvard College. All rights reserved.

Harvard authorizes you to view or download a single copy of the Harvard Content on EBSCOhost solely for your personal, noncommercial use if you include the following copyright notice:

"Copyright, President and Fellows of Harvard College. All rights reserved" and other copyright and proprietary rights notices which were contained in the Harvard Content. Reproduction and/or redistribution of the Harvard Content is expressly prohibited. Any special rules for the use of other items provided on EBSCOhost may be included elsewhere within the site and are incorporated into these Terms and Conditions.

The Harvard Content is protected by copyright under both United States and foreign laws. Title to the Harvard Content remains with President and Fellows, Harvard College. Any use of the Harvard Content not expressly permitted by these Terms and Conditions is a breach of these Terms and Conditions and may violate copyright, trademark, and other laws. Harvard Content and features are subject to change or termination without notice in the editorial discretion of Harvard. All rights not expressly granted herein are reserved to President and Fellows, Harvard College.

If you violate any of these Terms and Conditions, your permission to use the Harvard Content automatically terminates and you must immediately destroy any copies you have made of any portion of the Harvard Content.

MEDICAL DISCLAIMER

The information contained in this online site is intended to provide accurate and helpful health information for the general public. It is made available with the understanding that the author and publisher are not engaged in rendering medical, health, psychological, or any other kind of personal professional services on this site. The information should not be considered complete and does not cover all diseases, ailments, physical conditions or their treatment. It should not be used in place of a call or visit to a medical, health or other competent professional, who should be consulted before adopting any of the suggestions in this site or drawing inferences from it.

The information about drugs contained on this site is general in nature. It does not cover all possible uses, actions, precautions, side effects, or interactions of the medicines mentioned, nor is the information intended as medical advice for individual problems or for making an evaluation as to the risks and benefits of taking a particular drug.

The operator(s) of this site and the publisher specifically disclaim all responsibility for any liability, loss or risk, personal or otherwise, which is incurred as a consequence, directly or indirectly, of the use and application of any of the material on this site.

Post-Pancreatic Resection

General Overview and Unique Complications

Maureen Morrison, MSN, ACNP, DNPc

Pancreatic cancer is one of the most silently lethal forms of malignancy. With a lack of evidence-based research regarding pancreatic cancer, the evidence-based treatment options are limited. Currently, the best chance for a curative treatment is a pancreatic resection that typically is combined with chemotherapy and/or radiation therapy. The long-term treatment course and overall survival of the patient depend largely on the pathology report generated from the expurgated tissue analysis. After pancreatic resection, patients may experience both general and unique complications that, if not treated appropriately and assessed correctly, can be lethal. Provided here is a general overview of pancreatic resections and their unique postoperative complications for critical-care nurses. As the first-line provider, critical-care nurses assess and manage acute complications, in addition to educating patients to the long-term complications and how to manage them.

Keywords: Pancreatic cancer, Pancreatic resection, Postpancreatic resection

[DIMENS CRIT CARE NURS. 2010;29(4):157-162]

■ PANCREATIC CANCER

Pancreatic cancer is characterized by its very aggressive biological behavior. It can quickly become a rapidly disseminating and deadly tumor. Additionally, with pancreatic cancer's silent behavior, the diagnosis is often made after it is too late to pursue a curative treatment. The National Cancer Institute's Surveillance Epidemiological and End Results cancer statistics analysis reports that, in 2009 alone, there will be more than 40 470 people diagnosed with pancreatic cancer, whereas 35 240 people will succumb to the disease.¹ Of those diagnosed, only 7% of cases are diagnosed while the cancer is still locally confined at the primary site; 26% are diagnosed after the cancer has spread to regional lymph nodes or directly beyond the primary site; 53% are diagnosed after the cancer had already metastasized, and the remaining 14% of patients had unknown staging information.^{1,2}

The overall outcome of pancreatic cancer has not improved much over the last decade, with mortality continuing to soar above 95%.¹ Between 1998 and 2002, the median age at diagnosis for pancreatic cancer was 72 years, whereas the median age at death was 73 years.¹ The overall 5-year relative survival rate for a person diagnosed with pancreatic cancer between 1999 and 2005 was 5.5%. The corresponding 5-year relative survival rates were 22.2% for localized, 8.7% for regional, 1.8% for distant, and 4.9% for unstaged survivors.¹ Furthermore, upon surgery, approximately 1 in 10 patients considered preoperatively to have resectable tumors was found to have nonresectable disease intraoperatively. Pancreatic cancer is a very aggressive tumor, and although most diagnosed are treated by nonsurgical methods, the best chance for long-term survival is to pursue a potentially curative pancreatic resection. Because of the nature

Postpancreatic Resection

of the high degree of technical expertise required to perform a pancreatic resection, consultation at a specialized, high-volume pancreatic center will help to minimize operative mortality. Not only do post-pancreatic-resection patients require technically skilled surgeons, but they also depend greatly on the specially trained and educated critical-care nurses who treat them postoperatively.¹⁻³

The overall 5-year relative survival rate for a person diagnosed with pancreatic cancer between 1999 and 2005 was 5.5%.

If the patient who is about to undergo a pancreatic resection is correctly counseled prior to having an oper-

ation, he/she would have been told about the possibility of having to experience several postoperative complications (Table 1).³⁻¹³ However, it is important to keep in mind that, even with the most in-depth and detailed preoperative counseling, the patient is mostly likely extremely overwhelmed and will be in need of counseling more than once to allow the perioperative education time to be processed and absorbed. The patient is going to turn to the care providers whom he/she has first contact with after the operation, when the postoperative complications begin, for guidance in what to expect during the normal postoperative course. Therefore, critical-care nurses have become the first provider that the patients turn to for education, reassurance, and support.

■ THE PANCREAS

The pancreas is a comet-shaped organ, with a round head, a body, and a tapered tail. It sits between the spine

TABLE 1 Pancreatic Resection Postoperative Complications

Complications	Pathology	S/Sx	Treatment	Time of Onset and Duration
Gastric ileus	Gastrointestinal paralysis secondary to the trauma of an operation combined with delayed motility secondary to perioperative medications	Early satiety, postprandial nausea and vomiting, decreased bowel sounds, constipation, anorexia	Liquid diet, use of medications to stimulate gastric emptying, gastric decompression via gastrostomy tube release or possible nasogastric tube placement	POD 0-7; typically short term during initial healing of gastrointestinal trauma, but can continue with constipating pain medications
Endocrine insufficiency	If not enough healthy pancreatic tissue remains after resection, islet/acinar cell trauma, death, or removal	Hyperglycemia and associated symptoms	Exogenous insulin is typically required; it should be noted some oral antidiabetes medications can adversely affect the pancreas and should be avoided	POD 0-30; can be lifelong
Fistula	An abnormal connection between the pancreatic/bile ducts that cause the seepage of pancreatic and bilious fluids that can cause digestions of surrounding tissues in the peritoneal and abdominal cavity	Abdominal pain/distension, dullness to percussion, fever, chest pain, copious amounts of pancreatic enzyme/bilious fluid containing drainage	In acute situations, there is a possibility that the patient may need operative intervention, or possible insertion of, or reinsertion of, a percutaneous intra-abdominal drainage tube such as a Jackson-Pratt, combined with supportive therapies such as antipyretics	POD 0-7
Biliary reflux	The flow of bile upward from the small intestines to the stomach or esophagus due to biliary dyskinesia or biliary stricture	Chronic acid reflux, chronic nausea with bilious emesis, dyspepsia, halitosis, tooth enamel decay	Acid-suppressing medications and antiemetics	POD 0-45
Exocrine insufficiency	If not enough healthy pancreatic tissue remains after resection, islet/acinar cell trauma, death, or removal	Postprandial hyperperistalsis and diarrhea, malnutrition secondary to lack of fat absorption, weight loss	Exogenous pancreatic enzymes, anti-diarrhea medications, nutritional counseling and support	POD 0-30; can be lifelong

Abbreviations: POD, postoperative day; S/Sx, signs and symptoms.

and the stomach, with the head nestled in the curve of the duodenum, whereas the tail of the pancreas is tucked into the spleen. The pancreas is one of the major digestive and endocrine organs; it provides both endocrine and exocrine functions. The pancreas secretes digestive enzymes, such as amylase and lipase, into the small intestines from the main pancreatic duct where they help to digest carbohydrates, proteins, and fats to be later absorbed. The pancreas produces hormones such as insulin, glucagon, and somatostatin, which help regulate the endocrine system.^{14,15}

PANCREATIC RESECTIONS

There are several different types of pancreatic resections, and the type used is greatly dependent on the location of the mass that needs to be removed. Pancreatic resections vary widely; there is a total pancreatectomy, in addition to several other distinct subtotal pancreatectomies (Table 2).^{3-13,16-21} Total pancreatectomies are less common than subtotal resections as the postoperative complication of type 1 diabetes can seriously and permanently impact the quality of the patient’s life. This is particularly true as, according to the American Cancer Society, the average age of a person diagnosed with pancreatic cancer is 72 years, and the effect of diabetes in this population can be devastating on his/her overall health.^{2,6} The most common subtotal pancreatic resections include the pancreaticoduodenectomy and central or distal pancreatec-

tomies. Although each resection is unique in the technical aspects of the operation, several of the postoperative complications are similar (Tables 1 and 2). There are a few complications that are very unique to pancreatic resection, and those 3 are addressed in this article. The remaining complications are listed in Tables 1 and 2.

POSTOPERATIVE COMPLICATIONS

Pancreatic Fistula

Patients who have undergone pancreatic resection experience general postoperative complications such as hemorrhage and infection; however, several postoperative complications specific to pancreatic resection occur. Particularly, there is a tremendous amount of gastrointestinal and endocrine dysfunction including gastric ileus, pancreatic fistula, biliary reflux, pancreatic exocrine insufficiency, and pancreatic endocrine insufficiency (Table 2).

Several postoperative complications specific to pancreatic resection occur.

Pancreatic fistulas are one of the highest reported postoperative complications that occur after resection.¹⁸

TABLE 2 Pancreatic Resection Types and Associated Complications

Type of Resection	Common Postoperative Complications ^a
Total pancreatectomy: the radical resection of the pancreas	Lifelong pancreatic endocrine insufficiency and pancreatic exocrine insufficiency, biliary reflux
Pancreaticoduodenectomy: the diseased portion of the head of the pancreas is removed along with a segment of the duodenum, and a new connection between the two is made. Typically, this operation is part of what is known as the Whipple procedure	Short-term pancreatic endocrine/exocrine insufficiency, gastric ileus, biliary reflux, pancreatic fistula
Whipple procedure: removal of the diseased portion via a pancreaticoduodenectomy, antrectomy, cholecystectomy, choledochectomy with reconstruction of the anatomy via pancreaticojejunostomy, choledochojejunostomy, and a gastrojejunostomy	Lifelong pancreatic endocrine insufficiency and pancreatic exocrine insufficiency or short-term pancreatic endocrine/exocrine insufficiency, gastric ileus, biliary reflux, pancreatic fistula, postcholecystectomy syndrome
Central pancreatectomy: a middle or central portion of the pancreas is removed, preserving as much of the remaining tissue as possible	Short-term pancreatic endocrine/exocrine insufficiency, pancreatic fistula
Distal pancreatectomy: the distal portion of the pancreas is resected; typically, the spleen is also removed at this time, given probable involvement of the splenic artery	Short-term pancreatic endocrine/exocrine insufficiency, pancreatic fistula, asplenia
Pancreaticojejunostomy: a loop of the jejunum is mobilized and attached to the pancreatic duct to allow for better drainage of the pancreas	Short-term pancreatic endocrine/exocrine insufficiency, gastric ileus, biliary reflux, pancreatic fistula
Frey procedure: there is local resection at the head of the pancreas with a longitudinal pancreaticojejunostomy	Short-term pancreatic endocrine/exocrine insufficiency, gastric ileus, biliary reflux, pancreatic fistula, necrotic bowel syndrome
Beger procedure: in this pancreatic head resection, the duodenum is preserved	Short-term pancreatic endocrine/exocrine insufficiency, pancreatic fistula

^aExcludes general postoperative complications such as infection, DVT/PE, hemorrhage, etc.

Defined as the formation of an abnormal communication between the pancreas and other internal structures or organs that is developed secondary to surgical resection. A pancreatic fistula can be a difficult complication to manage and can occur both in the early and late stages of the postoperative healing course. If a fistula forms, it can allow the passage of digestive enzymes to flow into the peritoneal or pleural cavities, can cause pancreatic ascites, and eventually may track through the retroperitoneum into the mediastinum by way of the aortic or esophageal hiatus. If the secretions break through the mediastinal pleura, then a fistula into 1 or both pleural cavities can result. An internal pancreatic fistula should be suspected in any patient who presents with ascites or with a massive unilateral or bilateral pleural effusion. The diagnosis is confirmed by cross-sectional imaging, and then a therapeutic paracentesis or thoracentesis is performed. The fluid drained should be sent for analysis, the results of which will help to confirm diagnosis.^{8,18}

It is vital that these fistulas are diagnosed and treated early, given the ease with which these types of complications can progress. Critical-care nurses should be aware of the signs and symptoms of postoperative fistulas. Therefore, routine assessment for abdominal ascites or pulmonary dysfunction consistent with pleural effusions should be performed. Abdominal ascites is typically uncommon in a patient without a history of liver dysfunction, and its development should be followed up on immediately. Postoperative pleural effusions are considered more commonplace and may be associated with many different etiologies. Consideration of cross-sectional imaging to rule out a pancreatic fistula is appropriate in the setting of a persistent malingering or worsening effusion. Particular care should be used when evaluating for or treating pancreatic fistulas, given the limited and poor treatments available. Nurses administering care to patients after pancreatic resection should keep this complication, and the associated signs and symptoms, in mind during physical assessment.

Pancreatic Endocrine/Exocrine Insufficiency

When discussing pancreatic resection, it is important to remember the functions of the pancreas. The pancreas has two main functions: exocrine function and endocrine function. The pancreas has one main duct and several smaller tributary ducts that excrete digestive enzymes such as amylase and lipase into the duodenum of the small intestines. The enzymes mix with food to digest carbohydrates, proteins, and fats along the intestinal tract, so that the body is able to absorb that calories and nutrients it needs to maintain energy. This secretion of enzymes to digest food is the exocrine function of the pancreas. When the pancreas is damaged, or there is less

healthy tissue remaining to provide exocrine function, the patient experiences digestive dysfunction most commonly diagnosed by postprandial abdominal pain, diarrhea, and steatorrhea. Exocrine insufficiency can be treated by the intake of exogenous digestive enzymes a few minutes prior to eating, but if untreated both long- and short-term complications arise. In the short term, lack of nutritional absorption can cause poor wound healing, diarrhea-induced dehydration, electrolyte imbalance, and lethargy. Long-term exocrine insufficiency can lead to malnutrition and weight loss, and it is considered a poor prognostic indicator. A patient may develop pancreatic insufficiency as a result of intraoperative trauma to the pancreas, and as the pancreas heals, this complication may diminish; however, if there is not enough healthy pancreatic exocrine tissue remaining, this complication may be lifelong.^{9,11-13,21}

In addition to exocrine function, the pancreas is an endocrine gland. The pancreas produces hormones such as insulin and glucagon, in addition to other hormones, which have unique functions throughout the body. Particularly, the pancreas is known for its control of blood glucose levels, which is why, when the pancreas does not have enough healthy or functioning pancreatic endocrine cells remaining, the patient will develop post-pancreatic-resection insulin-dependent diabetes. It should be noted that almost all patients who undergo pancreatic resection will be maintained on exogenous insulin postoperatively for the first week as a result of pancreatic trauma. During the initial postoperative course, blood glucose levels are expected to be quite labile, as the pancreas may first function in short spurts during initial healing. Typically, by the fourth week postoperatively, the patient will know whether he/she will have permanent endocrine insufficiency. During the postoperative course, the patient will be responsible for learning how to check his/her blood glucose levels, so upon discharge home, he/she will be able to monitor and record his/her blood glucose levels. These levels are crucial for the first postoperative outpatient follow-up appointment, and patients typically require checking their blood glucose daily upon discharge home.^{11-13,22}

Critical-care nurses should be aware of the possible postoperative complications that can arise due to pancreatic endocrine or exocrine insufficiency. Nurses should be prepared to answer questions about the way pancreatic resection can affect the endocrine system. It should be noted that, given the average age of those diagnosed with pancreatic cancer is 72 years, the development of diabetes can be a difficult postoperative complication to learn how to manage. Diabetes education and training can seem daunting and be overwhelming for a patient to experience. The new diagnosis of diabetes can have a heavy emotional

impact, and although this complication can be transient postoperatively, all patients are at risk for developing permanent diabetes after pancreatic resection. It is in the patient's best interest to be aware of what diabetes is, and how it can affect his/her quality of life. The patient will need support when it comes to diabetes education and training, and the more the patient is exposed to diabetes training and education, the more reinforced the information becomes. Reinforcing this information provides the patient with the ability to take control and responsibility for keeping his/her blood glucose in check and at a safe level. Critical-care nurses should be prepared to facilitate the patient's diabetes training and education. Additionally, critical-care nurses should be evaluating the patient for diarrhea and postprandial pain that is associated with pancreatic exocrine insufficiency. Symptoms of either pancreatic exocrine or endocrine dysfunction should be evaluated during physical assessment.

Nurses should be prepared to answer questions about the way pancreatic resection can affect the endocrine system.

CONCLUSION

Pancreatic cancer is a very lethal malignancy that is not easily treated. The patient's limited treatment options include a major abdominal operation preceded and/or followed by oncologic therapies. These treatments are fraught with serious complications that can, if not managed appropriately, leave the patient with a very poor quality of life. To help the patient become his/her own best advocate, it is up to the providers to educate the patient about the complications and how to manage them effectively. As critical-care nurses are the frontline of providers caring for these patients, it is up to them to help the patient focus on what they should expect to experience and how to ensure they maintain a safe, positive treatment course.

Although pancreatic resections are not commonplace, particularly at smaller medical institutions, as technology improves and as more research on pancreatic cancer is conducted, there will continue to be an increasing number of patients requiring pancreatic resection. It is imperative that critical-care nurses are prepared to care for patients with pancreatic cancer who are status post pancreatic resection.^{12,13,22,23}

Critical-care nurses spend more time with the patient than any other multidisciplinary team member does during postoperative hospitalization. It is up to us to be knowledgeable about pancreatic cancer and the way

pancreatic resections affect our patients. By understanding the technical differences between resections, and the pathophysiology of how pancreatic resection affects the endocrine and gastrointestinal systems, assessment for unique complications is more effective. With effective assessment, early diagnosis and treatment of lethal complications can ensue, ultimately preventing mortality and securing improved overall survival, not to mention patient quality of life. We need to evaluate for those postoperative complications that are unique to pancreatic resection, and we need to help educate and inform our patients about what type of complications can arise, and how to be prepared to handle them. A nurse who is able to counsel a patient about what to expect after a pancreatic resection, that the complications are unique, but expected, and how the patient can prevent or handle these complications reduces the enormous amounts of fear and helplessness that the patient experiences.

Acknowledgment

The author thanks her colleagues at the Pancreas Center Columbia University Medical Center, New York.

References

1. Horner MJ, Ries LAG, Krapcho M, et al. *SEER Cancer Statistics Review, 1975-2006*. Bethesda, MD; 2008. National Cancer Institute. http://seer.cancer.gov/csr/1975_2006/. Based on November 2008 SEER data submission, posted to the SEER Web site.
2. The Pancreatic Cancer page. National Institute of Cancer Web site. <http://www.cancer.gov/cancertopics/types/pancreatic>. Accessed July 30, 2009.
3. The Bile duct cancer page. American Cancer Society Web site. http://www.cancer.org/docroot/CRI/content/CRI_2_4_4X_Surgery_69.asp?sitearea=. Accessed August 30, 2009.
4. Kimura W, Inoue T, Futakawa N, Shinkai H, Han I, Muto M. Spleen-preserving distal Pancreatectomy with conservation of the splenic artery and vein. *Surgery*. 1996;120(5):885-890. http://www.sciencedirect.com.monstera.cc.columbia.edu:2048/science?_ob=MIimg&_imagekey=B6WXC-4K3145B-J-1&_cdi=7155&_user=18704&_orig=browse&_coverDate=11%2F30%2F1996&_sk=998799994&view=c&wchp=dGLbVzzSkzS&md5=44ecf9634040b827ad406b541be9d40&ie=/sdarticle.pdf. Accessed June 10, 2009.
5. Mack LA, Kaklamanos IG, Ioannia G, et al. Gastric decompression and enteral feeding through a double-lumen gastrojejunostomy tube improves outcomes after pancreaticoduodenectomy. *Ann Surg*. 2004;240(5):845-851. <http://www.pubmedcentral.nih.gov.monstera.cc.columbia.edu:2048/articlerender.fcgi?artid=1356491>. Accessed July 30, 2009.
6. Allendorf John D, Schroppe Beth A, Inabnet William B, Chabot John A. Postoperative glycemic control in patients undergoing central pancreatectomy for mid-gland lesions. *World J Surg*. 2007;31:164-168.
7. Ocui LM, Sarmiento JM, Staley CA, et al. Comparison of central and extended left pancreatectomy for lesions of the pancreatic neck. *Ann Surg Oncol*. 2008;15(8):1534-4681. <http://www.springerlink.com.monstera.cc.columbia.edu:2048/content/4086t4rk8v3m5851/fulltext.pdf>. Accessed July 1, 2009.
8. Pratt W, Maithel AK, Vanounou T, Callery MP, Vollmer CM. Postoperative pancreatic fistulas are not equivalent after proximal, distal, and central pancreatectomy. *J Gastrointest Surg*. 2006;10(9):1264-1279. http://www.sciencedirect.com.monstera.cc.columbia.edu:2048/science?_ob=MIimg&_imagekey=B6WXC-4K3145B-J-1&_cdi=7155&_user=18704&_orig=browse&_coverDate=11%2F30%2F2006

- edu:2048/science?_ob=MImg&_imagekey=B6W8T-4MCBWT9-9-1&_cdi=6663&_user=18704&_orig=search&_coverDate=11%2F30%2F2006&_sk=999899990&view=c&wchp=dGLbVzW-zSkzV&cmd5=62dab4ce206a4a60f8bcb90c4f52065f&ie=/sdarticle.pdf. Accessed August 30, 2009.
9. Kahl S, Malfertheiner P. Exocrine and endocrine pancreatic insufficiency after pancreatic surgery. *Clin Gastroenterol.* 2004; 18(5):947-955.
 10. The Columbia University Medical Center. The Pancreas Center. <http://pancreasmd.org/>. Accessed June 10, 2009.
 11. Chaudhary A, Negi AA, Masood A, Thombare M. Complications after Frey's procedure for chronic pancreatitis. *Am J Surg.* 2004;188(3):277-281.
 12. Seiler CA, Wagner M, Bachman T, et al. Randomized clinical trial of pylorus-preserving duodenopancreatectomy versus classic Whipple resection—long term results. *Br J Surg.* 2005; 92(5):547-556. <http://www3.interscience.wiley.com.monstera.cc.columbia.edu:2048/cgi-bin/fulltext/110434045/PDFSTART?CRETRY=1&SRETRY=0>. Accessed August 25, 2009.
 13. Crippa S, Bassi C, Warshaw AL, et al. Middle pancreatectomy: indications, short- and long-term operative outcomes. *Ann Surg.* 2007;246(1):69-76. <http://www.pubmedcentral.nih.gov/picrender.fcgi?artid=1899210&blobtype=pdf>. Accessed August 25, 2009.
 14. Wilson JD, Foster DW. *William's Textbook of Endocrinology.* 8th ed. Philadelphia, PA: Harcourt Brace Javanovich, Inc; 1992.
 15. The Endocrine System page. Inner Body: Your Guide to Human Anatomy Online Web site. <http://www.innerbody.com/image/endoov.html>. Accessed June 10, 2008.
 16. Bassi C, Stocken DD, Olah A, et al. Influence of Surgical resection and post operative-complications on survival following adjuvant treatment for pancreatic cancer the ESPAC-1 randomized controlled trial. *Dig Surg.* 2005;22:353-363. <http://content.karger.com.monstera.cc.columbia.edu:2048/ProdukteDB/produkte.asp?Aktion=ShowPDF&ArtikelNr=000089771&Ausgabe=231418&ProduktNr=223996&filename=000089771.pdf>. Accessed July 17, 2009.
 17. Adham M, Giunipero A, Hervieu V, Courbiere M, Pattersky C. Central pancreatectomy. *Arch Surg.* 2008;143(2):175-181. <http://archsurg.ama-assn.org.monstera.cc.columbia.edu:2048/cgi/reprint/143/2/175>. Accessed July 14, 2009.
 18. Lipsett PA, Cameron JL. Internal pancreatic fistula. *Am J Surg.* 1992;163:216-220.
 19. Frey CF, Child CG, Fry W. Pancreatectomy for chronic pancreatitis. *Ann Surg.* 1978;403-412. <http://www.pubmedcentral.nih.gov/picrender.fcgi?artid=1345429&blobtype=pdf>. Accessed June 10, 2009.
 20. Gouma DJ, van Geenen RCI, van Gulik TM, et al. Rates of complications and death after pancreaticoduodenectomy: risk factors and the impact of hospital volume. *Ann Surg.* 2000; 232(6):786-795.
 21. Falconi M, Mantovani W, Crippa S, Mascetta G, Salvia R, Pederzoli P. Pancreatic insufficiency after different resections for benign tumours. *Br J Surg.* 2007;95(1):85-91. <http://www3.interscience.wiley.com.monstera.cc.columbia.edu:2048/cgi-bin/fulltext/117350580/PDFSTART>. Accessed August 22, 2009.
 22. Allendorf JD, Lauerma M, Bill A, et al. Neoadjuvant chemotherapy and radiation for patients with locally unresectable pancreatic adenocarcinoma: feasibility, efficacy, and survival. *J Gastrointest Surg.* 2008;12(1):91-100.
 23. Auter PK, Coleman J. Pancreatic cancer: a continuum of care. *Semin Oncol Nurs.* 1999;15(1):36-47.

ABOUT THE AUTHOR

Maureen Morrison, MSN, ACNP, DNPc, is an acute care nurse practitioner who is currently in her residency to complete the doctorate in clinical nursing at Columbia University School of Nursing, New York, New York.

Address correspondence and reprint request to: Maureen Morrison, MSN, ACNP, DNPc, 525 East 68th street, F-761, New York, NY 10065 (msm2131@columbia.edu).

**DCCN
on the Web**

Visit *Dimensions of Critical Care Nursing* on the Web. Take CE tests on-line, peruse back issues, order article reprints, or subscribe at <http://www.dccnjournal.com>.

Small Bowel Obstruction for Inpatients: Process

A CT of abdomen/pelvis with IV contrast is done to diagnose a small bowel obstruction. If diagnosed, the following process may be followed:

- Do not proceed with process if:
 - Infectious abdominal process
 - Cancer
 - Incarcerated hernia
 - Pregnant
 - Abdominal surgery within 6 weeks
 - History of pelvic radiation
 - Previous Roux-en-Y bypass
 - Concern for inguinal hernia
- ***Patients should be continuously assessed and re-evaluated for peritonitis throughout this process*** (increasing abdominal pain/tenderness/distention, fever, nausea/vomiting, loss of appetite, diarrhea, decreased urinary output, thirst, inability to pass stool or gas, fatigue)
 - Report to covering MD right away.
- Insert NGT
- Correct electrolytes and resuscitate with fluids (per MD orders)
 - Consult surgery if not already done
- Ensure NGT is patent, intra-gastric on x-ray, has been on suction for at least 2 hours
- Place head of bed > 30 degrees
- Administer **90 ml** of Gastrografin or Isovue 300 via NGT and **clamp for 1 hour then place back on suction** (Isovue 300 is the name of the contrast; 90 ml of either contrast is the correct dose to administer)
- Check portable KUB in 8 hours from contrast administration
 - If contrast ***not in cecum***, repeat KUB in 24 hours (if contrast still not in cecum, surgery should be considered)
 - If contrast ***is in cecum***, remove NGT and starts sips of liquid with advancement to full liquids within 24 hours

Contrast for CT Scan of Abdomen (Not for Small Bowel Obstruction)

- Contrast bottle comes from CT scan
- Directions are on the bottle (doses, timing of administration and when to send the patient to CT scan)
- Clamp NGT after administration of each dose, leave clamped through patient transport to radiology
- Contact radiology with any questions, ext. 44624
- Contrast is given in this case for diagnostic purposes, not therapeutic

AT THE BEDSIDE

Upper GI Bleeding

A 45-year-old white man is admitted with reports of an 18-hour history of nausea and vomiting of large amounts of "coffee-ground secretions" and frequent "maroon-colored" stools. He reports a previous history of peptic ulcer disease diagnosed at age 35. He has been hospitalized twice in the past for active GI bleeding. A duodenal ulcer near the pylorus on the posterior wall of the stomach was diagnosed by endoscopy. Significant findings on his admission profile were

Vital Signs

Blood pressure: 96/60 mm Hg lying;
90/58 mm Hg sitting
Heart rate: 120 beats/min; sinus
tachycardia
Respiratory rate: 32 breaths/min, deep
Temperature: 99.2°F (oral)

Respiratory

Breath sounds clear in all lung fields

Cardiovascular

S₁/S₂ no murmurs
Extremities cool, diaphoretic; pulses present but weak

Abdomen

Distended with hyperactive bowel sounds in all four quadrants

Tender right upper quadrant, no rebound tenderness

Neurologic

Alert, oriented
Anxious

Genitourinary

50 mL of amber, cloudy urine following Foley catheter insertion
Stools liquid maroon, guaiac positive

Arterial Blood Gases

pH	7.49
PaCO ₂	28 mm Hg
HCO ₃	19 mEq/L
PaO ₂	61 mm Hg on room air
SaO ₂	89%
Hematocrit	25%
Hemoglobin	7.0 g/dL
White blood cell count	17,000/mm ³
Prothrombin time	11 seconds
Activated partial thromboplastin time	30 seconds
Platelet count	110,000/mm ³
Serum potassium	3.5 mEq/L (decreased)
Serum sodium	150 mEq/L
Serum glucose	210 mg/dL
Serum blood urea nitrogen	40
Serum creatinine	0.9
Liver function	Within normal limits




Figure 14-3. Hypovolemic shock.




Figure 14-4. Upper GI bleeding treatment guide.